



IIT-JEE (MAIN/ADVANCED) • NEET • BOARDS • NTSE • KVPY






# CHEMISTRY QUESTION PAPER WITH SOLUTION

## (CODE – F3)




Address - Near Himfed Building, BCS Chowk, New Shimla (HP) - 171009  
[www.radianceiit.in](http://www.radianceiit.in) | Phone: +91 8091157087 | E-mail: [er.j.k.tiwari@gmail.com](mailto:er.j.k.tiwari@gmail.com)

46. Identify compound X in the following sequences of reactions :



Sol. 1



47. Identify a molecule which does not exist.

- (1) C<sub>2</sub>                      (2) O<sub>2</sub>                      (3) He<sub>2</sub>                      (4) Li<sub>2</sub>

Sol. 3


Diatomic molecule for which bond order is zero or negative does not exist.

Molecule	BO
C <sub>2</sub>	2
O <sub>2</sub>	2
He <sub>2</sub>	Zero
Li <sub>2</sub>	1

48. Which of the following is a natural polymer ?

- (1) polybutadiene                      (2) poly (Butadiene-acrylonitrile)  
 (3) cis-1,4-polyisoprene              (4) poly (Butadiene-styrene)

Sol. 3







49. An increase in the concentration of the reactants of a reaction leads to change in :

- (1) threshold energy (2) collision frequency  
 (3) activation energy (4) heat of reaction


Sol. 2

Collision frequency, as number of molecules per unit volume increases.

50. Anisole on cleavage with HI gives :




Sol. 3



51. The number of protons, neutrons and electrons in  $^{175}_{71}\text{Lu}$ , respectively, are :

- (1) 71, 71 and 104 (2) 175, 104 and 71  
 (3) 71, 104 and 71 (4) 104, 71 and 71

Sol. 3



no. of protons	=	71
no. of neutrons	=	$175 - 71 = 104$
no. of electrons	=	71

52. The calculated spin only magnetic moment of  $\text{Cr}^{2+}$  ion is :

- (1) 5.92 BM (2) 2.84 BM  
 (3) 3.87 BM (4) 4.90 BM



55. Match the following and identify the correct option.

- |   |   |
|---|---|
| (a) $\text{CO(g)} + \text{H}_2\text{(g)}$ | (i) $\text{Mg(HCO}_3)_2 + \text{Ca(HCO}_3)_2$ |
| (b) Temporary hardness of water           | (ii) An electron deficient hydride            |
| (c) $\text{B}_2\text{H}_6$                | (iii) Synthesis gas                           |
| (d) $\text{H}_2\text{O}_2$                | (iv) Non-planar structure                     |
- 
- |           |       |      |      |
|-----------|-------|------|------|
| (a)       | (b)   | (c)  | (d)  |
| (1) (iii) | (iv)  | (ii) | (i)  |
| (2) (i)   | (iii) | (ii) | (iv) |
| (3) (iii) | (i)   | (ii) | (iv) |
| (4) (iii) | (ii)  | (i)  | (iv) |

Sol. 3

$\text{CO(g)} + \text{H}_2\text{(g)} = \text{Water gas or synthesis gas}$

Temporary hardness of water is due to bicarbonates of  $\text{Ca}^{2+}$  &  $\text{Mg}^{2+}$

$\text{B}_2\text{H}_6$  is a electron deficient compound due to presence of banana bond.  $\text{H}_2\text{O}_2$  open book like structure which is non-planar

56. The mixture which shows positive deviation from Raoult's law is :

- |                          |                                |
|--------------------------|--------------------------------|
| (1) Acetone + Chloroform | (2) Chloroethane + Bromoethane |
| (3) Ethanol + Acetone    | (4) Benzene + Toluene          |

Sol. 3

Acetone + ethanol shows positive deviation from Raoult's law. Pure ethanol possesses H-bonding and adding acetone to ethanol causes breaking of some H-bonds. This causes increase in observed vapour pressure.

57. The freezing point depression constant ( $K_f$ ) of benzene is  $5.12 \text{ K kg mol}^{-1}$ . The freezing point depression for the solution of molality  $0.078 \text{ m}$  containing a non-electrolyte solute in benzene is (rounded off upto two decimal places) :

- |                      |                      |                      |                      |
|----------------------|----------------------|----------------------|----------------------|
| (1) $0.40 \text{ K}$ | (2) $0.60 \text{ K}$ | (3) $0.20 \text{ K}$ | (4) $0.80 \text{ K}$ |
|----------------------|----------------------|----------------------|----------------------|

Sol. 1


$T_f = i k_f m$

$$\begin{aligned} T_f &= 1 \times 5.12 \times 0.078 \\ &= 0.3993 \\ &= 0.40 \text{ K} \end{aligned}$$


58. Which of the following set of molecules will have zero dipole moment ?

- (1) Nitrogen trifluoride, beryllium difluoride, water, 1,3-dichlorobenzene
- (2) Boron trifluoride, beryllium difluoride, carbon dioxide, 1,4-dichlorobenzene
- (3) Ammonia, beryllium difluoride, water 1,4-dichlorobenzene
- (4) Boron trifluoride, hydrogen fluoride, carbon dioxide, 1,3-dichlorobenzene

**Sol. 2**




Non-polar molecule



Non-polar molecule

Non-polar molecule



Non-polar molecule

Cl

1,4-dichlorobenzene

**59.** A tertiary butyl carbocation is more stable than a secondary butyl carbocation because of which of the following ?

(1) -R effect of -CH<sub>3</sub> groups


(2) Hyperconjugation

(3) -I effect of -CH<sub>3</sub> groups

(4) + R effect of -CH<sub>3</sub> groups

**Sol. 2**

Tert butyl cation  
stable due to  
9  $\alpha$  - H Hyperconjugation



**60.** Find out the solubility of Ni(OH)<sub>2</sub> in 0.1 M NaOH. Given that the ionic product of Ni(OH)<sub>2</sub> is  $2 \times 10^{-15}$ .


(1)  $1 \times 10^{-13}$  M

(2)  $1 \times 10^{-8}$  M

(3)  $2 \times 10^{-13}$  M

(4)  $2 \times 10^{-8}$  M

**Sol. 3**



$$K_{\text{sp}} = [\text{Ni}^{2+}] [\text{OH}^-]^2$$

$$K_{\text{sp}} = s \times (0.1)^2$$

$$2 \times 10^{-15}$$

$$s = \frac{2 \times 10^{-15}}{0.1^2}$$

$$s = 2 \times 10^{-13} \text{ M}$$



**64.** A mixture of N<sub>2</sub> and Ar gases in a cylinder contains 7 g of N<sub>2</sub> and 8 g of Ar. If the total pressure of the mixture of the gases in the cylinder is 27 bar, the partial pressure of N<sub>2</sub> is:

[Use atomic masses (in g mol<sup>-1</sup>): N = 14, Ar = 40]

- (1) 15 bar                      (2) 18 bar                      (3) 9 bar                      (4) 12 bar

**Sol. 1**

According to Dalton's law

$$P_{N_2} = \frac{P_{N_2} \times \frac{7}{28}}{\frac{7}{28} + \frac{8}{40}} \times 27$$

$$= \frac{0.25}{0.2 + 0.25} \times 27$$

$$= 15 \text{ bar}$$

**65.** Which of the following is the correct order of increasing field strength of ligands to form coordination compounds ?

- (1) F<sup>-</sup> < SCN<sup>-</sup> < C<sub>2</sub>O<sub>4</sub><sup>2-</sup> < CN<sup>-</sup>                      (2) CN<sup>-</sup> < C<sub>2</sub>O<sub>4</sub><sup>2-</sup> < SCN<sup>-</sup> < F<sup>-</sup>  
 (3) SCN<sup>-</sup> < F<sup>-</sup> < C<sub>2</sub>O<sub>4</sub><sup>2-</sup> < CN<sup>-</sup>                      (4) SCN<sup>-</sup> < F<sup>-</sup> < CN<sup>-</sup> < C<sub>2</sub>O<sub>4</sub><sup>2-</sup>

**Sol. 3**

Fact from spectrochemical series

**66.** Paper chromatography is an example of :

- (1) Thin layer chromatography                      (2) Column chromatography  
 (3) Adsorption chromatography                      (4) Partition chromatography


**Sol. 4**

It is an example of partition chromatography as mixture of substance are separated by partition.

**67.** Sucrose on hydrolysis gives:

- (1) -D-Glucose + -D-Fructose                      (2) -D-Fructose + -D-Fructose  
 (3) -D-Glucose + -D-Fructose                      (4) -D-Glucose + -D-Glucose

**Sol. 1**





68. The rate constant for a first order reaction is  $4.606 \times 10^{-3} \text{ s}^{-1}$ . The time required to reduce 2.0 g of the reactant to 0.2 g is:

- (1) 500 s                      (2) 1000 s                      (3) 100 s                      (4) 200 s

Sol. 1

$$t = \frac{1}{k} \ln \frac{C_0}{C_t}$$

$$= \frac{1}{4.606 \times 10^{-3}} 2.303 \log \frac{2}{0.2}$$


$$= \frac{1}{2 \times 10^{-3}} \log 10$$

$$= 500 \text{ sec}$$

69. Reaction between benzaldehyde and acetophenone in presence of dilute NaOH is known as:

- (1) Cross Cannizzaro's reaction                      (2) Cross Aldol condensation  
 (3) Aldol condensation                      (4) Cannizzaro's reaction

Sol. 2



Two different carbonyl compound is treated with base known as crossed aldol condensation.

70. Which of the following is not correct about carbon monoxide?

- (1) The carboxyhaemoglobin (haemoglobin bound to CO) is less stable than oxyhaemoglobin.  
 (2) It is produced due to incomplete combustion.  
 (3) It forms carboxyhaemoglobin.  
 (4) It reduces oxygen carrying ability of blood.

Sol. 1

CO when combine with haemoglobin it forms carboxyhaemoglobin which more stable than oxyhaemoglobin thats why it reduces oxygen carring ability of blood.

71. Hydrolysis of sucrose is given by the following reaction



If the equilibrium constant ( $K_c$ ) is  $2 \times 10^{13}$  at 300 K, the value of  $rG^\circ$  at the same temperature will be:

- (1)  $8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(3 \times 10^{13})$   
(2)  $-8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(4 \times 10^{13})$   
(3)  $-8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(2 \times 10^{13})$   
(4)  $8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 300 \text{ K} \times \ln(2 \times 10^{13})$

Sol. 3

$$G^\circ = -RT \ln K$$

$$G^\circ = -8.314 \times 300 \times \ln(2 \times 10^{13})$$

72. HCl was passed through a solution of  $\text{CaCl}_2$ ,  $\text{MgCl}_2$  and  $\text{NaCl}$ . Which of the following compounds(s) crystallise(s)?

- (1) only  $\text{MgCl}_2$  (2)  $\text{NaCl}$ ,  $\text{MgCl}_2$  and  $\text{CaCl}_2$   
(3) Both  $\text{MgCl}_2$  and  $\text{CaCl}_2$  (4) Only  $\text{NaCl}$

Sol. 1

$\text{MgCl}_2$  has least solubility among  $\text{CaCl}_2$ ,  $\text{MgCl}_2$  &  $\text{NaCl}$  that's why it crystallises first on passing HCl.

73. An element has a body centered cubic (bcc) structure with a cell edge of 288 pm. The atomic radius is:

- (1)  $\frac{4}{\sqrt{3}} \times 288 \text{ pm}$  (2)  $\frac{4}{\sqrt{2}} \times 288 \text{ pm}$  (3)  $\frac{\sqrt{3}}{4} \times 288 \text{ pm}$  (4)  $\frac{\sqrt{2}}{4} \times 288 \text{ pm}$

Sol. 3

For bcc


$$\sqrt{3}a = 4r$$




$$r = \frac{\sqrt{3}}{4} \times 288 \text{ pm}$$

74. Which of the following oxoacid of sulphur has -O-O- linkage?

- (1)  $\text{H}_2\text{S}_2\text{O}_8$ , peroxodisulphuric acid (2)  $\text{H}_2\text{S}_2\text{O}_7$ , pyrosulphuric acid  
(3)  $\text{H}_2\text{SO}_3$ , sulphurous acid (4)  $\text{H}_2\text{SO}_4$ , sulphuric acid

Sol. 1





**75.** Identify the incorrect statement.

(1) Interstitial compounds are those that are formed when small atoms like H, C or N are trapped inside the crystal lattices of metals

(2) The oxidation states of chromium in  $\text{CrO}_4^{2-}$  and  $\text{Cr}_2\text{O}_7^{2-}$  are not the same

(3)  $\text{Cr}^{2+}(\text{d}^4)$  is a stronger reducing agent than  $\text{Fe}^{2+}(\text{d}^6)$  in water

(4) The transition metals and their compounds are known for their catalytic activity due to their ability to adopt multiple oxidation states and to form

complexes **Sol. 2**

(1) Interstitial compounds are those compounds that are formed when small atoms like H, B, C or N are trapped inside crystal lattices of metals.

(2) Oxidation state of chromium in  $\text{CrO}_4^{2-}$  &  $\text{Cr}_2\text{O}_7^{2-}$  are +6 (same)

(3) Fact

$\text{Cr}^{2+}$  is stronger reducing agent than  $\text{Fe}^{2+}$  in water on the basis of standard reduction potential.

(4) Transition metals are one of the best available catalyst due to their ability to adopt multiple oxidation states & to form complexes.

**76.** Which of the following is a cationic detergent?

(1) Cetyltrimethyl ammonium bromide      (2) Sodium dodecylbenzene sulphonate

(3) Sodium lauryl sulphate                      (4) Sodium stearate

**Sol. 1**

Cetyltrimethyl ammonium bromide is a cationic detergent as its hydrophilic part is a quaternary ammonium ion.

77. The correct option for free expansion of an ideal gas under adiabatic condition is:

- (1)  $q < 0$ ,  $T = 0$  and  $w = 0$                       (2)  $q > 0$ ,  $T > 0$  and  $w > 0$   
(3)  $q = 0$ ,  $T = 0$  and  $w = 0$                       (4)  $q = 0$ ,  $T < 0$  and  $w > 0$

Sol. 3

For an adiabatic process,  $q = 0$

for free expansion,  $P_{\text{ext}} = 0$

$$w = 0$$

& from First law of Thermodynamics,  $E = 0$

or  $T = 0$

78. On electrolysis of dil. sulphuric acid using platinum (Pt) electrode, the product obtained at anode will be:

- (1)  $\text{H}_2\text{S}$  gas                      (2)  $\text{SO}_2$  gas                      (3) Hydrogen gas                      (4) Oxygen gas

Sol. 4

Anode :  $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$

Cathode :  $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$

oxygen gas will be obtained at anode.

79. Identify the correct statement from the following:

- (1) Vapour phase refining is carried out for Nickel by Van Arkel method  
(2) Pig iron can be moulded into a variety of shapes  
(3) Wrought iron is impure iron with 4% carbon  
(4) Blister copper has blistered appearance due to evolution of  $\text{CO}_2$

Sol. 2

(1) Vapour phase refining is carried out for nickel by Mond process.

(2) Pig iron is molten iron obtained from blast furnace after smelting it can be moulded into variety of shapes.


(3) Wrought iron is the purest form of iron which has less than 4% carbon.

(4) Blister copper has blistered appearance due to evolution of  $\text{SO}_2$ .

80. Which of the following is a basic amino acid?

- (1) Tyrosine                      (2) Lysine                      (3) Serine                      (4) Alanine

Sol. 2



Basic part is more than acidic part so lysine is basic amino acid.

81. Identify the incorrect match.

Name	IUPAC Official Name
(a) Unnilunium	(i) Mendelvium
(b) Unniltrium	(ii) Lawrencium
(c) Unnilhexium	(iii) Seaborgium
(d) Unununnium	(iv) Darmstadtium
(1) (c), (iii)	(2) (d), (iv)
(3) (a), (i)	(4) (b), (ii)

Sol. 2

101 - Unnilunium - Mendeleevium  
103 - Unniltrium - Lawrencium  
106 - Unnilhexium - Seaborgium  
111 - Unununium - Roentgenium

82. Which of the following alkane cannot be made in good yield by Wurtz reaction?

(1) n-Heptane      (2) n-Butane      (3) n-Hexane      (4) 2,3,-Dimethylbutane


Sol. 1

Wurtz reaction is used to prepare symmetrical alkane.

83. Elimination reaction of 2-Bromo-pentane to form pent-2-ene is:

(a) -Elimination reaction      (b) Follows Zaitsev rule  
(c) Dehydrohalogenation reaction      (d) Dehydration reaction  
(1) (b), (c), (d)      (2) (a), (b), (d)      (3) (a), (b), (c)      (4) (a), (c), (d)

Sol. 3



84. The number of Faradays (F) required to produce 20g of calcium from molten  $\text{CaCl}_2$  (Atomic mass of Ca =  $40 \text{ g mol}^{-1}$ ) is:

(1) 3      (2) 4      (3) 1      (4) 2

Sol. 3

$\text{Equivalent}_{\text{Ca}} = \text{Equivalent}_{\text{current}}$

$\frac{W}{MM} \cdot nf = \text{No. of Faradays}$

$\text{No. of Faradays} = 40 \frac{20}{2}$

$\text{No. of Faradays} = 1$



85. Which one of the followings has maximum number of atoms?

- (1) 1g of O<sub>2</sub>(g) (Atomic mass of O=16)
- (2) 1g of Li(s) (Atomic mass of Li=7)
- (3) 1g of Ag(s) (Atomic mass of Ag = 108)
- (4) 1g of Mg(s) (Atomic mass of Mg = 24)

Sol. 2

$$(1) \text{ no. of atoms} = \frac{1}{7} N_A$$

$$(2) \text{ no. of atoms} = \frac{1}{108} N_A$$

$$(3) \text{ no. of atoms} = \frac{1}{24} N_A$$


$$(4) \text{ no. of atoms} = \frac{1}{32} \times 2 N_A$$

1 gm of Li is having maximum no. of atoms.

86. For the reaction, 2Cl(g) → Cl<sub>2</sub>(g), the correct option is:

- (1) ΔH < 0 and ΔS > 0
- (2) ΔH < 0 and ΔS < 0
- (3) ΔH > 0 and ΔS > 0
- (4) ΔH > 0 and ΔS < 0

Sol. 2



Due to bond formation stability increases which results in release of heat.

H<sub>r</sub> = -ve or exothermic process

$$H_r < 0$$

ΔS < 0, because number of Cl atoms decreases in the formation of Cl<sub>2</sub>(g)

87. Identify the correct statements from the following:

- (a) CO<sub>2</sub>(g) is used as refrigerant for ice cream and frozen food
- (b) The structure of C<sub>60</sub> contains twelve six carbon rings and twenty five carbon rings.
- (c) ZSM-5, a type of zeolite, is used to convert alcohols into gasoline
- (d) CO is colourless and odourless gas

- (1) (b) and (c) only
- (2) (c) and (d) only
- (3) (a), (b) and (c) only
- (4) (a) and (c) only

Sol. 2

CO<sub>2</sub> (s) is used as refrigerant for ice cream & frozen food. six carbon rings – 20

C<sub>60</sub> — five carbon rings – 12

ZSM-5 zeolite used to convert alcohols into gasoline

CO is colourless & odourless gas

- 88.** Measuring Zeta potential useful in determining which property of colloidal solution?  
(1) Stability of the colloidal particles      (2) Size of the colloidal particles  
(3) Viscosity      (4) Solubility


**Sol. 1**

A large +ve or -ve value of Zeta potential indicate good stability

- 89.** What is the change in oxidation number of carbon in the following reaction?  $\text{CH}_4(\text{g}) + 4\text{Cl}_2(\text{g}) \rightarrow \text{CCl}_4(\text{l}) + 4\text{HCl}(\text{g})$

(1) -4 to +4      (2) 0 to -4      (3) +4 to +4      (4) 0 to +4

**Sol. 1**



- 90.** The following metal ion activates many enzymes, participates in the oxidation of glucose to produce ATP and with Na, is responsible for the transmission of nerve signals.

(1) Calcium      (2) Potassium      (3) Iron      (4) Copper

**Sol. 2**

Potassium